

Northumbria Research Link

Citation: Ludwig, Björn, Hourfar, Jan, Glasl, Bettina and Ludwig, Gregory (2013) Relocating canines into the dental arch: the role of self-litigating brackets with additional slots. Orthodontic Products. pp. 76-83. ISSN 1097-797X

Published by: Novicom

URL: <http://www.orthodonticproductsonline.com/orp-clini...>
<<http://www.orthodonticproductsonline.com/orp-clinical-tips/15549-relocating-canines-into-the-dental-arch>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/11690/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Northumbria Research Link

Citation: Ludwig, B., Hourfar, J., Glasl, B. and Ludwig, G. (2013) Ludwig, B., Hourfar, J., Glasl, B. & Ludwig, G. (2013)"Relocating canines into the dental arch: the role of self-litigating brackets with additional slots", Orthodontic Products, pp.76-83. Orthodontic Products. pp. 76-83. ISSN 1097-797X

Published by: Torrance, CA : Novicom, Inc.,

URL: <http://www.orthodonticproductsonline.com>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/11690/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

www.northumbria.ac.uk/nrl



Relocating Canines into the Dental Arch

BY BJÖRN LUDWIG, DMD, MSD; JAN HOURFAR, DMD, MSD; BETTINA GLASL, DMD, MSD; AND GREGORY LUDWIG, PhD, MBA, BA (HONS), BBA

Apart from mandibular third molars, maxillary canines are most frequently affected by dislocation, triggering the need for therapeutic treatment. Not including wisdom teeth, maxillary canines represent about 60% of teeth selected for relocation. Among the pathological causes are shortage of space and eruption problems, as well as an atypical germ situation. Furthermore, genetical causes are often discussed. Particularly relevant in this context is the so-called guidance theory because it emphasizes the role of lateral incisors as a central stabilizing factor that allows for alignment of canine teeth, which in turn is of high importance for other teeth.¹⁻²

Physiological eruption of upper teeth usually takes place at a physical age of about 11.5 years.³ In the majority of cases, this happens on both sides at about the same time. Around 10% of cases, however, reveal a difference of up to 1 year in direct comparison of time of eruption at both sides. The prevalence rate of dislocation of maxillary canines is about 0.8% to 2.3%, with the rate of female cases slightly higher.⁴ Moreover, 8% of patients are affected by bilateral relocation. About one-third of retained canine teeth are located in a labial position within the alveolar process, and about two-thirds are located in palatal positions.⁵

Clinically, dislocation of canine teeth is often easy to identify even without medical imaging because vestibular canines may be palpated unilaterally. Further indicators could be peg-shaped lateral incisors or their aplasia. The latter case imposes an obvious excess of space.

X-ray radiographs providing an overview of the patient's teeth highlight a generally better prognosis for successful integration

if the angle between teeth axis and occlusal plane is higher than 45°. In turn, successful orthodontic integration of dislocated canines allows preservation of functionally very important canines.

Practical Considerations

Prior to any relocation of canine teeth, a surgical cut-down is usually required. This can be implemented via open or closed



BJÖRN LUDWIG, DMD, MSD, is in private practice in Traben-Trarbach, Germany. He is scientific coordinator at the University Homburg/Saar. He can be reached at bludwig@kieferorthopaedie-mosel.de.



JAN HOURFAR, DMD, MSD, is in a private practice in Reinheim, Germany. He can be reached at hourfar@t-online.de.



BETTINA GLASL, DMD, MSD, has a private practice together with Björn Ludwig in Traben-Trarbach, Germany. She can be reached at bglasl@kieferorthopaedie-mosel.de.



GREGORY LUDWIG, PhD, MBA, BA (Hons), BBA, is senior lecturer in the faculty of business and law at Northumbria University in the United Kingdom. He can be reached at g.ludwig@northumbria.ac.uk.

cloud9ortho

"it just makes sense"

Stop By
Booth 325
at AAO
for a chance to WIN
Visa Gift Cards & a Surface Pro

Run Your Practice Without the Worry & Cost of Servers

NO Servers NO T1 Lines Lower IT Costs

- No software to install
- Use on Macs or PCs with no emulators
- Images load fast and are available instantly
- Satellite offices run just as fast as main location

"I've got two practices in two states, one practice with Cloud9 and one without. Needless to say, I wish they were BOTH Cloud9. I've had so many of my colleagues tell me that their practice management system is "on the cloud", which simply isn't true. With Cloud9 I have no servers, no worries about backups and no worries about the latest update costing me more money. I travel a lot, and frequently need access to patient images and charts. Terminal services is archaic and painful. I never have remote log in headaches with Cloud9, no matter the computer or location, I'm always in my practice. Don't be fooled into thinking you're on the cloud unless your software was built from the cloud up."



- Dr. John Graham

"I'm scrambling to get ready for a lecture next week. I gotta pull together tons of case photos and have NO TIME to do it between now and then. I turned on my "Personal HotSpot" and connected my laptop. BAM I have access to Cloud9 and all my patient photos and charts. So I am sitting here making my PowerPoint and keep bringing up Cloud9 and snipping and copying photos right into my PowerPoint. So yup....access from hotels, the beach, the golf course, airports, even airplanes. Anytime---Anywhere!"

- Dr. Doug Depew



Schedule your Demo Today!
1-800-394-6050

info@cloud9ortho.com



www.cloud9ortho.com



Figure 1: (A) Forestadent BioQuick bracket with auxiliary slot (dimension .016" x .016").

techniques, depending on localization of the respective tooth. Prompt active elongation is recommended in the postoperative period. The process of orthodontic gap opening has often been initiated at this point in time. Deep intra-osseous dislocation often necessitates prior extrusion; this could be realized using ballista springs,⁶ for example. Following sufficient extrusion of the teeth (aided by respective appliances), it can then be relocated into the dental arch.

In theory, orthodontists can choose from a variety of options. However, some of these offer striking disadvantages. If the relocation "pulls" via an elastic main arch, strong reciprocal side effects are to be expected, mainly affecting teeth neighboring the gap. This shortcoming particularly applies in cases displaying strongly ectopic canines. On the other hand, if flexible sutures leading from ectopic teeth toward the rigid orthodontic arch are selected, poor esthetics and hygiene are often the result.

Furthermore, a considerable decrease in pulling power can be predicted in such cases.

Another option is a flexible parallel arch, in addition to the rigid orthodontic main arch within the bracket slot. The main disadvantage in this case is the friction and barrier to teeth movement, because the parallel arch has restricted movement. Reciprocal side effects, meanwhile, can have a negative impact on overall treatment success if, for example, palatal canines are buccally shifted via the extension of a quad helix. This is due to general difficulties in controlling shifting powers of quad helix structures. Too strong, nonphysiological pulling power may result in hyalinization and, in some cases, even total suspension of teeth movement. Root ankylosis is very rare in maxillary canines.⁷

Advantages of SL Bracket Usage with Auxiliary Slots

Self-litigating brackets with additional, auxiliary slots⁸ offer a good alternative to the above-discussed treatment options. Not all SL systems, however, are suitable. Inserting of additional parallel arches is often not feasible and would immediately result in dysfunctional sealing caps. Forestadent's Quick bracket system addresses this issue through a 0.016" x 0.016" auxiliary slot (Figure 1). This auxiliary slot thus aids any treatment and relocation of dislocated or ectopical canines. In addition, it has a number of major advantages. This is because it allows for insertion of parallel arches (eg, 0.012" NiTi) that can move freely almost without any friction, in cases of rigid orthodontic main arches in situ. In turn, this allows for the release of continuous physiological pulling power, effectively preventing potential hyalinization and respectively longer treatment periods. Throughout the physiologically aided integration of canine teeth, further physical teeth movements may then be realized at the rigid orthodontic main arch, offering a major contribution toward achieving overall treatment goals.

Clinical Examples

In Figures 2 and 3, we see the vestibular auxiliary slot technique post-laser (Sirona) gingivectomy. We used a Forestadent BioQuick bracket with auxiliary slot (dimension .016" x .016"). In Figure 2, we see the left vestibular retained canine within the attached gingiva following laser gingivectomy. The main arch was .019" x .025" SS and was combined with an overlay arch (.012" BioStarter) within the auxiliary slot.



Figure 2: Left vestibular retained canine within the attached gingiva following laser gingivectomy; main arch .019" x .025" SS and overlay arch (.012" BioStarter) within auxiliary slot.



Figure 3: Gingiva position post-treatment. A circumscriptive gingivoplasty is planned for the integrated canine in order to improve aesthetic results.

AN EARTH-SHAKING PRODUCT SET TO TURN THE ORTHODONTIC WORLD ON ITS SIDE



**SEE WHAT'S SHAKING
AT THE AAO ANNUAL SESSION!**

Ormco Booth #1009

ormco.com/braceyourself





Figure 4: Vestibular auxiliary slot technique with closed exposure. (A) CBCT-data set in 3D volume image: high left vestibular canine - dislocated. (B) High vestibular canine - dislocated; main arch .019" x .025" SS and overlay arch (.012" BioStarter) within auxiliary slot. (C) The canine tooth receives a +17° palatal torque bracket in the finishing phase. This is required to be able to move the canine tooth root into the palatal bone for achievement of healthy periodontal conditions.




Figure 5: Vestibular auxiliary slot technique with closed exposure. (A) Vestibular highly dislocated canine tooth; main arch .019" x .025" SS and overlay arch (.012" BioStarter) within auxiliary slot. (B) Gingiva condition post-treatment completion. The gingival situation could have been further improved through more palatal torque at the canine tooth.



We´ll be in the spotlight!

**You're invited to take part on May 4TH
at our AAO Annual Session Booth #1901.**



Figure 6: Palatal auxiliary slot technique. (A) CBCT data set in 3D imaging (upper jaw): considerable palatal dislocation of canine teeth. (B) Palatal relocated canine teeth post-laser exposure and adhesive attachment of catching chain; main arch .019" x .025" SS and overlay arch (.012" BioStarter) within auxiliary slot.

Meanwhile, Figure 4 (page 80) shows a vestibular auxiliary slot technique with closed exposure. The CBCT data set in 3D volume image shows a dislocation of a high left vestibular canine (Figure 4A, page 80). Again the main arch was .019" x .025" SS, and the overlay arch (.012" BioStarter) was within the auxiliary slot (Figure 4B, page 80). In the

final stage, the canine tooth received a +17° palatal torque bracket. This is required to be able to move the canine tooth root into the palatal bone to achieve healthy periodontal conditions (Figure 4C, page 80).

Figure 5 (page 80) shows another vestibular auxiliary slot technique with closed exposure. In this case, the vestibular canine was

dislocated. The main arch was .019" x .025" SS and the overlay arch (.012" BioStarter) within the auxiliary slot. Post-treatment, the gingival situation could have been further improved through more palatal torque at the canine tooth, but the x-ray post-treatment completion showed satisfying results in relation to the canine position and root alignment.

Infalab



Figure 4: (C) Natural gingiva position post-active treatment.

Finally, Figure 6 shows once more the palatal auxiliary slot technique. The CBCT data set in 3D imaging (upper jaw) shows a considerable palatal dislocation of the canine. The canine was exposed with a laser followed by adhesive attachment of a catching chain. The main arch was .019" x .025" SS, and the overlay arch (.012" BioStarter) was

within the auxiliary slot. Post-active treatment, the gingival margin looks natural (Figure 7).

Conclusion

Overall, employment of SL brackets with additional auxiliary slots offers a number of advantages for integration of dislocated or ectopically positioned canines: relatively short

treatment times due to easy operation and handling, uncomplicated insertion of parallel arches into the auxiliary slots, and decreased friction and physiological impact. **OP**



References for this article are available with **Orthodontic Products'** interactive and online editions.



DX Retainer & Splint

Tru-Tain offers the DX Retainer & Splint Material, a blend of medical-grade, FDA-approved plastics that combine durability, impact, strength, and wear resistance. The material is easy to form and bonds with acrylic. It may be used for retainers, minor tooth movement, and bite ramps. It is stronger than the original Tru-Tain retainer and splint plastic. Available in .030 and .040 thickness. 25, 50 and 100 sheet packs. Square 5" x 5", or round 125mm and 120mm. Contact the company for free samples.

For more information, contact

Tru-Tain

2625 Hwy 14 W,
Rochester, MN 55901;
(800) 843-0904; www.tru-tain.com



ARCH-101

architects : consultants : advocates



ARCH-101 provides architectural, planning, and consulting services for success-minded owners of specialty dental and healthcare practices.

Staunch advocates of our client's goals, we provide the highest-functioning solutions through principal-managed projects, design excellence, and client education.

arch-101.com

411 E Coeur d'Alene Ave Ste 101

Coeur d'Alene, ID 83814

(208) 391-7100

Scott Somers, NCARB, Architect

ssomers@arch-101.com